

# $\mathbb{Z}_p, \mathbb{Q}_p,$ AND THE RING OF WITT VECTORS

YOSHIFUMI TSUCHIMOTO

No.03:  $\boxed{\mathbb{Z}_p \text{ as a projective limit of } \{\mathbb{Z}/p^k\mathbb{Z}\}}$

DEFINITION 3.1. An ordered set  $\Lambda$  is said to be **directed** if for all  $i, j \in \Lambda$  there exists  $k \in \Lambda$  such that  $i \leq k$  and  $j \leq k$ .

DEFINITION 3.2. Let  $\Lambda$  be a directed set. Let  $\{X_\lambda\}_{\lambda \in \Lambda}$  be a family of topological rings. Assume we are given for each pair of elements  $(\lambda, \mu) \in \Lambda^2$  such that  $\lambda \geq \mu$ , a continuous homomorphisms

$$\phi_{\mu\lambda} : X_\lambda \rightarrow X_\mu.$$

We say that such a system  $(\{X_\lambda\}, \{\phi_{\mu\lambda}\})$  is a **projective system** of topological rings if it satisfies the following axioms.

- (1)  $\phi_{\nu\mu}\phi_{\mu\lambda} = \phi_{\nu\lambda}$  ( $\forall \lambda, \forall \mu \forall \nu$  such that  $\lambda \geq \mu \geq \nu$ ).
- (2)  $\phi_{\lambda\lambda} = \text{id}$  ( $\forall \lambda \in \Lambda$ ).

DEFINITION 3.3. Let  $\mathcal{X} = (\{X_\lambda\}, \{\phi_{\mu\lambda}\})$  be a projective system of topological rings. Then we say that a **projective limit**  $(X, \{\phi_\lambda\})$  of  $\mathcal{X}$  is given if

- (1)  $X$  is a topological ring.
- (2)  $\phi_\lambda : X \rightarrow X_\lambda$  is a continuous homomorphism.
- (3)  $\phi_{\mu\lambda} \circ \phi_\lambda = \phi_\mu$  for  $\forall \mu, \lambda$  such that  $\lambda \geq \mu$ .
- (4)  $(X, \{\phi_\lambda\})$  is a universal object among objects which satisfy (1)-(3).

The “universal” here means the following: If  $(Y, \psi_\lambda)$  satisfies

- (1)  $Y$  is a topological ring.
- (2)  $\psi_\lambda : Y \rightarrow X_\lambda$  is a continuous homomorphism.
- (3)  $\phi_{\mu\lambda} \circ \psi_\lambda = \psi_\mu$  for  $\forall \mu, \lambda$  such that  $\lambda \geq \mu$ .

Then there exists a unique continuous homomorphism

$$\Phi : Y \rightarrow X$$

such that

$$\psi_\lambda = \phi_\lambda \circ \Phi (\forall \lambda \in \Lambda).$$

PROPOSITION 3.4. *For any projective system of topological rings, a projective limit of the system exists. It is unique up to a unique isomorphism. (Hence we may call it **the** projective limit of the system.)*

DEFINITION 3.5. For any projective system  $(X, \{\phi_\lambda\})$  of topological rings, We denote the projective limit of it by

$$\varprojlim_{\lambda} X_{\lambda}.$$

Note: projective limits of systems of topological spaces, rings, groups, modules, and so on, are defined in a similar manner.

THEOREM 3.6.

$$\mathbb{Z}_p \cong \varprojlim_{k \rightarrow \infty} (\mathbb{Z}/p^k\mathbb{Z})$$

as a topological ring.

COROLLARY 3.7.  $\mathbb{Z}_p$  is a compact space.

Note: There are several ways to prove the result of the above corollary. For example, the ring  $\mathbb{Z}$  with the metric  $d_p$  is easily shown to be totally bounded.

PROPOSITION 3.8. Each element of  $\mathbb{Z}_p$  is expressed uniquely as

$$[0.a_1a_2a_3a_4\dots]_p \quad (a_i \in \{0, 1, \dots, p-1\} \quad (i = 1, 2, 3, \dots)).$$

EXERCISE 3.1. Is  $-4 = 1 - 5$  invertible in  $\mathbb{Z}_5$ ? (Hint: use formal expansion

$$(1 - x)^{-1} = 1 + x + x^2 + \dots$$

is it possible to write down a correct proof to see that the result is true?)