## COHOMOLOGIES.

## YOSHIFUMI TSUCHIMOTO

## 02. "Hom" modules.

LEMMA 2.1. Let R be a ring. Let  $f: M \to N$  be a homomorphism of R-modules. Then for any R-module L we may define:

- (1) A homomorphism  $\operatorname{Hom}_R(L,f): \operatorname{Hom}_R(L,M) \to \operatorname{Hom}_R(L,N)$  defined by  $\operatorname{Hom}_R(L,f)(g) = f \circ g$ .
- (2) A homomorphism  $\operatorname{Hom}_R(f,L) : \operatorname{Hom}_R(N,L) \to \operatorname{Hom}_R(M,L)$  defined by  $\operatorname{Hom}_R(f,L)(h) = h \circ f$ .

Proposition 2.2. Let R be a ring. Let

$$0 \to M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \to 0$$

be an exact sequence of R-modules. Then for any R-module N, we have:

(1)

- $0 \to \operatorname{Hom}_{R}(N, M_{1}) \stackrel{\operatorname{Hom}_{R}(N, f)}{\to} \operatorname{Hom}_{R}(N, M_{2}) \stackrel{\operatorname{Hom}_{R}(N, g)}{\to} \operatorname{Hom}_{R}(N, M_{3})$ is exact. The third arrow  $\operatorname{Hom}_{R}(N, g)$  need not be surjective.
  (2)
- $0 \to \operatorname{Hom}_R(M_3, N) \overset{\operatorname{Hom}_R(g, N)}{\to} \operatorname{Hom}_R(M_2, N) \overset{\operatorname{Hom}_R(f, N)}{\to} \operatorname{Hom}_R(M_1, N)$  is exact. The third arrow  $\operatorname{Hom}_R(f, N)$  need not be surjective.

EXERCISE 2.1. We consider an exact sequence

$$0 \to 3\mathbb{Z} \xrightarrow{i} \mathbb{Z} \to \mathbb{Z}/3\mathbb{Z} \to 0$$

where i is the inclusion map. Show that

$$\operatorname{Hom}_{\mathbb{Z}}(\mathbb{Z},\mathbb{Z}) \stackrel{\operatorname{Hom}(i,\mathbb{Z})}{\to} \operatorname{Hom}_{\mathbb{Z}}(3\mathbb{Z},\mathbb{Z})$$

is not surjective

EXERCISE 2.2. Assume R is a field. Then show that the third arrow which appear in the sequence (1) in Proposition 2.2 is surjective.