

A note on quasi-polarized manifolds (X, L) with $$h^0(K_X + (n - 2)L) = 0$$

YOSHIAKI FUKUMA

September 14, 2011

Let X be a projective variety of dimension n over the field of complex numbers \mathbb{C} , and let L be a nef and big (resp. an ample) line bundle on X . Then we call the pair (X, L) a *quasi-polarized* (resp. *polarized*) *variety*, and (X, L) is called a *quasi-polarized* (resp. *polarized*) *manifold* if X is smooth.

In this note, we consider a characterization of quasi-polarized manifolds (X, L) such that L is generated by its global sections and $h^0(K_X + (n - 2)L) = 0$.

Definition 1 Let (X, L) be a quasi-polarized variety of dimension n . Then the Euler-Poincaré characteristic $\chi(tL)$ is a polynomial in t of total degree n . We set

$$\chi(tL) = \sum_{j=0}^n \chi_j(L) \binom{t+j-1}{j}.$$

Here we will define the i th sectional geometric genus of quasi-polarized varieties.

Definition 2 ([2, Definition 2.1]) Let (X, L) be a quasi-polarized variety of dimension n and let i be an integer with $0 \leq i \leq n$. Then the i th *sectional geometric genus* $g_i(X, L)$ is defined by the following.

$$g_i(X, L) = (-1)^i (\chi_{n-i}(L) - \chi(\mathcal{O}_X)) + \sum_{j=0}^{n-i} (-1)^{n-i-j} h^{n-j}(\mathcal{O}_X).$$

Remark 1 (1) If $i = 0$, then $g_0(X, L) = L^n$.

(2) If $i = 1$, then $g_1(X, L)$ is the sectional genus $g(X, L)$ of (X, L) .

(3) If $i = n$, then $g_n(X, L) = h^n(\mathcal{O}_X)$.

Then by [2, Corollary 3.4] we get the following.

Theorem 1 Let (X, L) be a quasi-polarized manifold of dimension $n \geq 3$. Assume that L is spanned by its global sections. Then $h^0(K_X + (n - 2)L) = 0$ if and only if $g_2(X, L) = h^2(\mathcal{O}_X)$.

Moreover if L is ample and spanned by its global sections, then we get the following by [2, Corollary 3.5].

Theorem 2 Let (X, L) be a polarized manifold of dimension $n \geq 3$. Assume that L is spanned by its global sections. Then $g_2(X, L) = h^2(\mathcal{O}_X)$ if and only if (X, L) is one of the following types.

(1) $(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$.

- (2) $(\mathbb{Q}^n, \mathcal{O}_{\mathbb{Q}^n}(1))$.
- (3) A scroll over a smooth projective curve.
- (4) $K_X \sim -(n-1)L$, that is, (X, L) is a Del Pezzo manifold.
- (5) A hyperquadric fibration over a smooth curve.
- (6) $(\mathbb{P}_S(\mathcal{E}), H(\mathcal{E}))$, where S is a smooth projective surface and \mathcal{E} is an ample vector bundle of rank $n-1$ on S .
- (7) Let (M, A) be a reduction of (X, L) .
 - (7.1) $n = 4$, $(M, A) = (\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(2))$.
 - (7.2) $n = 3$, $(M, A) = (\mathbb{Q}^3, \mathcal{O}_{\mathbb{Q}^3}(2))$.
 - (7.3) $n = 3$, $(M, A) = (\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3}(3))$.
 - (7.4) $n = 3$, M is a \mathbb{P}^2 -bundle over a smooth curve C and $(F', A|_{F'}) \cong (\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$ for any fiber F' of it.

By Theorems 1 and 2, we get a characterization of polarized manifolds (X, L) with $h^0(K_X + (n-2)L) = 0$.

Next we consider the case where L is nef and big and spanned by its global sections. Here we use notation and terminologies in [1].

Theorem 3 *Let (X, L) be a quasi-polarized manifold of dimension $n \geq 3$. Assume that L is spanned by its global sections. Then $h^0(K_X + (n-2)L) = 0$ if and only if first reduction (X'', L'') of (X, L) in the sense of [1, Definition 5.5] satisfies one of the pairs in [1, Proposition 3.5, B)].*

Proof. (1) First we consider the case where (X, L) satisfies $h^0(K_X + (n-2)L) = 0$. Assume that $K_X + (n-2)L$ is pseudoeffective. Since L is spanned by its global sections, there exist $(n-3)$ members Y_1, \dots, Y_{n-3} of $|L|$ such that $X_{n-3} := Y_1 \cap \dots \cap Y_{n-3}$ is smooth. Let $L_{n-3} := L|_{X_{n-3}}$. Then we note that $K_{X_{n-3}} + L_{n-3}$ is pseudoeffective. Hence by [1, Theorem 5.6, 2)] $K_{X''_{n-3}} + L''_{n-3}$ is nef, where (X''_{n-3}, L''_{n-3}) is a first reduction of (X_{n-3}, L_{n-3}) in the sense of [1, Definition 5.5]. Since X''_{n-3} has only \mathbb{Q} -factorial terminal singularities, [3, Theorem 1.5] implies that $0 < h^0(K_{X''_{n-3}} + L''_{n-3}) = h^0(K_{X_{n-3}} + L_{n-3})$. Moreover we can easily check that $h^0(K_{X_{n-3}} + L_{n-3}) \leq h^0(K_X + (n-2)L)$. But this contradicts the assumption that $h^0(K_X + (n-2)L) = 0$. Hence $K_X + (n-2)L$ is not pseudoeffective. Therefore by [1, Theorem 5.6, 1)] first reduction (X'', L'') of (X, L) satisfies one of the pairs in [1, Proposition 3.5, B)].

(2) Next we consider the case where first reduction (X'', L'') of (X, L) satisfies one of the pairs in [1, Proposition 3.5, B)]. Then by [1, Theorem 5.6, 1)] we have $h^0(K_X + (n-2)L) = 0$. This completes the proof. \square

By Theorems 1 and 3, we also get the following.

Theorem 4 *Let (X, L) be a quasi-polarized manifold of dimension $n \geq 3$. Assume that L is spanned by its global sections. Then $g_2(X, L) = h^2(\mathcal{O}_X)$ if and only if first reduction (X'', L'') of (X, L) in the sense of [1, Definition 5.5] satisfies one of the pairs in [1, Proposition 3.5, B)].*

Remark 2 There exists a typo in the last case in [1, Proposition 3.5, B)].

Error	Correction
\dots is $\mathcal{O}(1)$.	\dots is $\mathcal{O}(2)$.

References

- [1] M. Andreatta, *Minimal model program with scaling and adjunction theory*, arXiv:1107.4878v1.
- [2] Y. Fukuma, *On the sectional geometric genus of quasi-polarized varieties, I*, *Comm. Algebra* 32 (2004), 1069–1100.
- [3] A. H\"oring, *On a conjecture of Beltrametti and Sommese*, arXiv:0912.1295, to appear in *J. Algebraic Geom.*

Yoshiaki Fukuma
Department of Mathematics
Faculty of Science
Kochi University
Akebono-cho, Kochi 780-8520
Japan
E-mail: fukuma@kochi-u.ac.jp